

NEW ZEALAND CONSUMER MEDICINE INFORMATION

GOSERELIN (TEVA)

Goserelin, 10.8 mg implant, in a prefilled syringe

What is in this leaflet

This leaflet answers some common questions about Goserelin (Teva) 10.8 mg. It does not contain all the available information. It does not take the place of talking to your doctor or pharmacist.

All medicines have risks and benefits. Your doctor will have weighed the risks of you receiving Goserelin (Teva) 10.8 mg against the benefits they expect it will have for you.

If you have any concerns about using this medicine, ask your doctor or pharmacist.

Keep this leaflet with the medicine. You may need to read it again.

What Goserelin (Teva) is used for

Goserelin (Teva) 10.8 mg is a special preparation of goserelin acetate which is designed to work in the body for 12 weeks (3 months) after each injection. Most people will need one injection for each 12 weeks (3 months) of treatment. Goserelin (Teva) 10.8 mg is otherwise similar to Goserelin (Teva) 3.6 mg which you may have received in the past. Goserelin (Teva) 3.6 mg lasts for only 28 days.

Goserelin (Teva) 10.8 mg is a member of the anti-hormonal group of medicines. This means that it affects the levels of various hormones (natural chemicals produced by the body). In men it will reduce levels of the male hormone, testosterone. In women it will reduce the levels of the female hormone, oestrogen.

In women, female hormones make the breasts grow, prepare the womb and other sex organs for pregnancy, and increase the sex drive.

In men, male hormones make the testicles grow, assist an erection, and increase the sex drive. When Goserelin (Teva) 10.8 mg lowers sex hormones, all these actions will be reduced. Due to the action of Goserelin (Teva) 10.8 mg on hormones, Goserelin (Teva) 10.8 mg can treat very different illnesses in both men and women.

MEN

Goserelin (Teva) 10.8 mg can treat prostate cancer in some men. It is not a cure for prostate cancer and does not work in every patient with prostate cancer.

WOMEN

Goserelin (Teva) 10.8 mg may be used for endometriosis instead of surgery. Goserelin (Teva) 10.8 mg is not a cure for endometriosis and may be helpful by reducing:

- “patches” of endometriosis tissue in the abdomen and pelvic area
- symptoms of endometriosis, including pain.

Endometriosis has no connection with any type of cancer.

Goserelin (Teva) 10.8 mg may be used for uterine fibroids before surgery. Fibroids are growths in the womb which are not cancer. Goserelin (Teva) 10.8 mg is not a cure for uterine fibroids and may be helpful by reducing the size of fibroids in or around the womb and symptoms of fibroids including pain.

Women who are receiving Goserelin (Teva) 10.8 mg for endometriosis or fibroids are not being treated for cancer. If you want more information about this ask your doctor or pharmacist.

Goserelin (Teva) 10.8 mg is not addictive.

Goserelin (Teva) 10.8 mg is only available with a doctor's prescription.

Before you are given Goserelin (Teva)

Before you start to use it

You must tell your doctor if:

- **you intend to become pregnant or plan to breast feed.**

Do not use Goserelin (Teva) 10.8 mg if you are pregnant, are trying to get pregnant, or while breastfeeding.

- **you have risk factors for loss of bone calcium and osteoporosis.**

These factors include:

- smoking
- having a slight build
- having a diet low in calcium
- having treatment with steroids for other conditions
- poor mobility (for example, confined to a wheelchair)

When women are treated with Goserelin (Teva) 10.8 mg, the amount of calcium in their bones decreases over a period of months. Some recovery of this loss can occur when treatment has stopped. A similar condition also happens after the change of life, when the natural production of female sex hormones falls. The link between the loss of calcium while on Goserelin (Teva) 10.8 mg and the loss of calcium after the change of life (menopause) is uncertain. However, women who have a family history of osteoporosis and other risk factors should discuss the use of Goserelin (Teva) 10.8 mg with their doctor before using Goserelin (Teva) 10.8 mg.

- **you have had an allergy to any other medicines or any other substances, such as preservatives or dyes.**
- **you have had any problems passing urine or if you have had any lower back problems.**
- **you have diabetes.**
- **you have any of the following: heart or blood vessel conditions, including heart rhythm problems (arrhythmia), or are being treated with medicines for these conditions.**

The risk of you having further heart rhythm problems may increase if you are taking Goserelin (Teva) 10.8 mg.

If you have not told your doctor about any of the above, tell him/her before you receive any Goserelin (Teva) 10.8 mg.

Tell your doctor if you are taking any other medicines, including medicines that you buy without a prescription from your pharmacy, supermarket or health food shop.

Goserelin (Teva) 10.8 mg might interfere with some medicines used to treat heart rhythm problems or might increase the risk of heart rhythm problems when used with some other medicines that can cause heart rhythm abnormalities.

When you must not use it

Do not use Goserelin (Teva) 10.8 mg if:

- **you are pregnant, could be pregnant or plan to get pregnant** unless you are receiving it as part of a treatment for infertility or have been specially advised by your doctor.
- **you are breastfeeding.** If you are breast feeding you should discuss this with your doctor.
- you have ever had an allergic reaction to Goserelin (Teva) 10.8 mg.
- the packaging is torn or shows signs of tampering.
- **the expiry date on the pack has passed.** If you receive this medicine after the expiry date has passed, it may not work.

Goserelin (Teva) 10.8 mg is not to be used in children.

How Goserelin (Teva) is given

How much you will be given

Goserelin (Teva) 10.8 mg will be given to you as an injection by your doctor or nurse. Goserelin (Teva) 10.8 mg comes in a special syringe and needle. A small pellet containing Goserelin (Teva) 10.8 mg is injected under the skin.

The injection is often at the front of the stomach (belly), or to the side of the stomach.

The injection leaves a small pellet under your skin. This pellet is designed to give you a dose of Goserelin (Teva) 10.8 mg which is released over 12 week (3 months). This means your injection will be given every 12 weeks (3 months).

How long to use it

Your doctor will tell you how long you may need to receive Goserelin (Teva) 10.8 mg. It is important that you carry on receiving your Goserelin (Teva) 10.8 mg even if feeling well unless your doctor decides it is time to stop treatment.

Women should only receive Goserelin (Teva) 10.8 mg for a maximum of 6 months.

If you miss a treatment

You should not delay or miss your 12 weekly (3 monthly) Goserelin (Teva) 10.8 mg. If you have missed a dose on purpose because you do not like a side effect of the medicine discuss this with your doctor. Talk to your doctor if you do not want to receive any more Goserelin (Teva) 10.8 mg because of side effects or for any other reason.

While you are using Goserelin (Teva)

Things you must do

Make sure to keep your appointments for your Goserelin (Teva) 10.8 mg dose every 3 months (12 weeks).

If you are a woman receiving therapy with Goserelin (Teva) 10.8 mg, barrier methods of contraception such as the condom or diaphragm (cap) should be used.

Oral forms of contraception (the "Pill") should not be taken when receiving Goserelin (Teva) 10.8 mg. **If you become pregnant while receiving Goserelin (Teva) 10.8 mg, tell your doctor immediately.**

If you go into hospital, let the medical staff know you are receiving Goserelin (Teva) 10.8 mg.

If you are about to be started on any new medicine, tell your doctor or pharmacist that you are receiving Goserelin (Teva) 10.8 mg.

You should only stop receiving Goserelin (Teva) 10.8 mg if advised to do so by your doctor.

It is unlikely that Goserelin (Teva) 10.8 mg will affect your ability to drive a car or to operate machinery.

If you are unsure about any of these points or have further questions please tell your doctor or pharmacist.

Side effects

Do not be alarmed by the following list of possible side effects. You may not experience any of them. Goserelin (Teva) 10.8 mg begins to work in the two weeks after the first injection. You should expect to notice changes in your body because your sex hormones will fall to low levels.

Tell your doctor or pharmacist as soon as possible if you do not feel well while you are using Goserelin (Teva) 10.8 mg.

Goserelin (Teva) 10.8 mg helps most people with endometriosis, fibroids, or prostate cancer, but it may have unwanted side effects in a few people. All medicines can have side effects. Sometimes they are serious, most of the time they are not. You may need medical treatment if you get some of the side effects.

Ask your doctor or pharmacist to answer any questions you have.

Tell your doctor if you notice any of the following and they worry you.

- rare allergic reactions
- skin rashes
- painful joints
- tingling in fingers or toes
- changes in blood pressure
- thinning of bones

Pain, bruising, bleeding, redness, swelling and other reactions may occur around the site of injection.

Injection site injury (including damage to blood vessels in the abdomen) has been reported following injection of Goserelin (Teva) 10.8 mg. In very rare cases this has caused severe bleeding. Contact your doctor immediately if you experience any of the following symptoms: abdominal pain, abdominal distension, shortness of breath, dizziness, low blood pressure and/or altered levels of consciousness.

In the very rare case, when patients receiving Goserelin (Teva) 10.8 mg have a tumour in their pituitary gland, Goserelin (Teva) 10.8 mg may make the tumour bleed or collapse. This can cause severe headaches, sickness, loss of eyesight and unconsciousness.

These are side effects in both men and women.

In males, sometimes side effects may be symptoms of the prostate cancer you are being treated for. Therefore you should always tell your doctor if you notice any possible side effect.

You may have a reduced sex drive with Goserelin (Teva) 10.8 mg. You may not be able to get an erection or father a child. However, Goserelin (Teva) 10.8 mg is not a contraceptive for men and should never be used that way.

You may get hot flushes, sweating and weight gain. Some men may get swollen or tender breasts. The testicles may become smaller. Mood swings have been commonly reported.

When you first start receiving Goserelin (Teva) 10.8 mg you may feel some pain in your bones. If this happens tell your doctor and you may be given something for this.

Very occasionally you may have trouble passing urine or experience lower back pain. If this happens, tell your doctor and you may be given something for this.

In females, most women get signs of the change of life (menopause) while on Goserelin (Teva) 10.8 mg such as:

- change in libido (sex drive)
- hot flushes and sweating
- mood changes, including depression
- a change of breast size
- headache
- a dry vagina
- weight gain
- hair loss
- acne

Formation of ovarian cysts may occur, which may result in pain in some women.

Goserelin (Teva) 10.8 mg will usually stop your periods. Some women will have a menstrual period in the first 2 weeks after the first injection of Goserelin (Teva) 10.8 mg. Rarely, some women may enter their natural menopause when being treated with Goserelin (Teva) 10.8 mg and will not resume having periods when the treatment with Goserelin (Teva) 10.8 mg stops. In some women, the time it takes for periods to begin again after they stop receiving Goserelin (Teva) 10.8 mg, can be prolonged.

Vaginal bleeding may occur. If you have fibroids, a slight increase in symptoms such as pain may occur. These effects are usually short-lived and discontinue on continuation of treatment. If symptoms persist or you are uncomfortable, contact your doctor. In addition, if you experience excessive nausea, vomiting or thirst, you should tell your doctor. This may indicate

possible changes in the amount of calcium in your blood and your doctor may have to do certain blood tests.

Important: This leaflet alerts you to some of the situations when you should call your doctor. Other situations, which cannot be predicted, may arise. Nothing in this leaflet should stop you from calling your doctor or pharmacist with any questions or concerns you have about using Goserelin (Teva) 10.8 mg.

In case of overdose

Getting too much Goserelin (Teva) 10.8 mg is unlikely as overdose is getting more than one Goserelin (Teva) 10.8 mg treatment in 12 weeks (3 months).

If a person received more than one injection in 12 weeks (3 months), Goserelin (Teva) 10.8 mg would act for a bit longer. Tell your doctor if you think you have received too much Goserelin (Teva) 10.8 mg.

Telephone your doctor or the National Poisons Information Centre Ph: 0800 POISON (0800 764 766) or go to Accident & Emergency at your nearest hospital immediately if you think that you or anyone else may have taken too much Goserelin (Teva) 10.8 mg. Even if there are no signs of discomfort or poisoning.

Storage conditions

Storage

Keep Goserelin (Teva) 10.8 mg the original packet. Let only your doctor or nurse open the pack.

Keep in a cool dry place, protected from light, where temperature stays below 30°C. Do not store it or any other medicine in the bathroom or near a sink. Do not leave it in the car or on window sills. Heat and dampness can destroy some medicines.

Keep it where children cannot reach it. A locked cupboard at least one and a half metres above the ground is a good place to store medicines.

Disposal

Once your doctor has given you Goserelin (Teva) 10.8 mg, he or she will dispose of the syringe and needles. If your Goserelin (Teva) 10.8 mg has passed its expiry date, ask your doctor or pharmacist how to dispose of it.

Product Description

What Goserelin (Teva) looks like

Goserelin (Teva) 10.8 mg comes in a special syringe and needle. The small pellet containing Goserelin (Teva) 10.8 mg is about the size of a grain of rice.

The pellet located inside the syringe. Goserelin (Teva) 10.8 mg is available in a pack of one injection.

Ingredients

Goserelin (Teva) 10.8 mg implant contains 10.8 mg of the active ingredient, goserelin (as goserelin acetate).

Goserelin (Teva) 10.8 mg also contains the following inactive ingredients: Poly(D,L-lactide), Poly(D,L-lactide-co-glycolide) 75:25.

Who supplies this medicine

Teva Pharma (New Zealand) Limited
PO Box 128 244
Remuera
Auckland 1541
Telephone: 0800 800 097

Date of information

16 November 2020