

# Medicianz Mupirocin Ointment



## *Mupirocin (2% w/w)*

---

### New Zealand Consumer Medicine Information

---

#### ***What is in this leaflet***

Please read this leaflet carefully before you use Medicianz MUPIROCIN Ointment.

This leaflet answers some common questions about Medicianz MUPIROCIN Ointment. It does not contain all the available information

It does not take the place of talking to your doctor or pharmacist.

All medicines have benefits and risks. Your doctor has weighed the risks of you using the Medicianz MUPIROCIN Ointment against the benefits they expect it will have for you.

If you are helping someone else to use Medicianz MUPIROCIN Ointment, read this leaflet before applying the first application.

**If you have any concerns about being treated with Medicianz MUPIROCIN Ointment, ask your doctor or pharmacist.**

This leaflet was last updated on the date at the end of this leaflet. More recent information may be available. The latest Consumer Medicine Information is available from your pharmacist or doctor and

may contain important information about the medicine and its use of which you should be aware.

**Keep this leaflet with the medicine.**

You may need to read it again.

---

#### ***What Medicianz MUPIROCIN Ointment is used for***

Medicianz MUPIROCIN Ointment contains mupirocin (2% w/w) as the active ingredient.

MUPIROCIN belongs to a group of medicines called antibiotics.

Antibiotics work by killing bacteria which can cause infection.

Medicianz MUPIROCIN Ointment is used for the treatment of certain types of skin infections such as school sores (impetigo), infection of the hair root and boils. Medicianz MUPIROCIN Ointment is also used when infection occurs in skin damaged by eczema, dermatitis, psoriasis, herpes (cold sores), wounds, cuts, grazes, insect bites, minor burns etc.

Medicianz MUPIROCIN Ointment is for use on your skin only.

**Ask your doctor if you have any questions about why this medicine has been prescribed for you.**

Your doctor may have prescribed Medicianz MUPIROCIN Ointment for another reason.

Medicianz MUPIROCIN Ointment is not addictive.

---

#### ***Before you use Medicianz MUPIROCIN Ointment***

##### ***When you should not be given it***

You must not be given Medicianz MUPIROCIN Ointment if you have an allergy to:

- any medicine containing mupirocin
- any of the ingredients listed at the end of this leaflet.

Some of the symptoms of an allergic reaction may include:

- shortness of breath
- wheezing or difficulty breathing

- swelling of the face, lips, tongue or other parts of the body
- rash, itching or hives on the skin.

**Do not use Medicianz MUPIROCIN Ointment after the expiry date printed on the pack, or the ointment shows visible signs of deterioration, or if the packaging is torn or shows signs of tampering.**

If it has expired or is damaged, return it to your pharmacist for disposal.

**If you are not sure whether you should start using this medicine, talk to your doctor.**

### ***Before you start to use it***

**Tell your doctor if you have allergies to any other medicines, foods, preservatives or dyes.**

**Tell your doctor if you are breastfeeding, pregnant or trying to become pregnant.**

Your doctor will discuss the risks and benefits with you.

**If you are using Medicianz MUPIROCIN Ointment on a cracked nipple, wash the nipple well before breast feeding your baby.**

**Medicianz MUPIROCIN Ointment should not be used to treat infections in or near the eyes, nostrils (in babies) or mouth.**

If Medicianz MUPIROCIN Ointment is accidentally

applied to the eyes, wash the eyes thoroughly with water.

**Medicianz MUPIROCIN Ointment contains polyethylene glycol which can be absorbed by the body from open wounds and damaged skin.**

**Polyethyleneglycol is removed from the body by the kidneys. Do not use Medicianz MUPIROCIN Ointment on large areas of broken or damaged skin, especially if you suffer from reduced kidney function.**

**Medicianz MUPIROCIN Ointment should not be used to treat areas of skin near tubes inserted into your body for the delivery or removal of fluid (cannulae). Medicianz MUPIROCIN Ointment should not be used in combination with cannulae.**

**If you develop an allergic reaction (such as a rash, local pain or swelling), wipe off the ointment, stop using Medicianz MUPIROCIN Ointment and tell your doctor.**

**Medicines like Medicianz MUPIROCIN Ointment can cause inflammation of the colon (large intestine), causing diarrhoea, usually with blood and mucus, stomach pain, fever (pseudomembranous colitis).**

**Tell your doctor as soon as possible if you get any of these symptoms.**

**If you have not told your doctor about any of the above, tell him/ her before**

**you start using Medicianz MUPIROCIN Ointment.**

### ***Taking other medicines***

**Tell your doctor if you are taking any other medicines, including any that you get without a prescription from your pharmacy, supermarket or health food shop.**

**Do not mix Medicianz MUPIROCIN Ointment with any other preparation. They may prevent Medicianz MUPIROCIN Ointment from working.**

---

### ***How to use Medicianz MUPIROCIN Ointment***

---

#### ***How much to use***

**Follow all directions given to you by your doctor or pharmacist carefully.**

They may differ from the information contained in this leaflet.

The usual recommendation is for a small amount to be applied to the affected area three times a daily.

If you have any questions about Medicianz MUPIROCIN Ointment and how to use it, please ask your doctor or pharmacist for help.

#### ***How to use it***

**Always wash your hands before and after applying Medicianz MUPIROCIN Ointment.**

With a piece of clean cotton wool or a gauze swab, apply a small quantity of Medicianz

MUPIROCIN Ointment to cover the affected area.

After applying the ointment, you may cover the treated area with a sterile bandage or gauze dressing, unless your doctor has told you to leave it uncovered.

### ***How long to use it***

Use Medicianz MUPIROCIN Ointment for as long as your doctor or pharmacist tells you to. Medicianz MUPIROCIN Ointment is usually applied for up to 10 days.

**It is important that you take the full course of Medicianz MUPIROCIN Ointment. Do not stop early as your symptoms may disappear before the infection is fully cleared.**

### ***If you forget to use it***

If you forget an application, apply some Medicianz MUPIROCIN Ointment as soon as you can, then continue as before.

### ***Use in children***

The use of Medicianz MUPIROCIN Ointment in children under 4 weeks of age is not recommended.

### ***If you use too much (overdose)***

Immediately telephone your doctor or National Poisons Centre (telephone 0800 POISON or 0800 764 766) for advice, if you think you or anyone else may have swallowed a large quantity of Medicianz MUPIROCIN Ointment, even if there are

**no signs of discomfort or poisoning.**

You may need urgent medical attention.

**If you are not sure what to do, contact your doctor, pharmacist or nearest hospital.**

---

### ***While you are using Medicianz MUPIROCIN Ointment***

---

#### ***Things you must do:***

**If you are about to be started on any new medicine, remind your doctor and pharmacist that you are using Medicianz MUPIROCIN Ointment.**

**Tell any other doctors, dentists, and pharmacists who treat you that you have been given this medicine.**

**If you become pregnant while using this medicine, tell your doctor immediately.**

**Tell your doctor if, for any reason, you have not used your medicine exactly as directed.**

Otherwise, your doctor may think that it was not working as it should and change your treatment unnecessarily.

#### ***Things you must not do***

**Do not give this medicine to anyone else, even if their symptoms seem similar to yours.**

**Do not use Medicianz MUPIROCIN Ointment to treat any other complaints unless your doctor says to.**

Do not stop using Medicianz MUPIROCIN Ointment, or change the dose without first checking with your doctor.

---

### ***Side effects***

---

**Tell your doctor or pharmacist as soon as possible if you do not feel well while you are using Medicianz MUPIROCIN Ointment.**

This medicine helps most people, but it may have unwanted side effects in a few people. All medicines can have side effects. Sometimes they are serious, most of the time they are not. You may need medical attention if you get some of the side effects.

**Check with your doctor as soon as possible if you think you are experiencing any side effects or allergic reactions due to taking Medicianz MUPIROCIN Ointment, even if the problem is not listed below.**

**Do not be alarmed by the following lists of side effects. You may not experience any of them.**

The most commonly reported side effects are:

- Itching
- Dryness
- Burning
- Redness
- Stinging
- Pain
- Swelling

Allergic reactions such as a rash, hives or swelling can occur with the use of

Medicianz MUPIROCIN  
Ointment.

**If you get a skin reaction,  
stop using Medicianz  
MUPIROCIN ointment.  
Wipe off any cream and tell  
your doctor as soon as  
possible.**

**Tell your doctor if you notice  
any of the following:**

- Swelling, itching, burning or pain in the area where Medicianz MUPIROCIN Ointment is applied.
- Wheezing, swelling of the lips/ mouth, difficulty in breathing, hayfever, lumpy rash (hives) or fainting. These could be symptoms of an allergic reaction.
- Collapse or loss of consciousness.

This is not a complete list of all possible side-effects. Others may occur in some people and there may be some side-effects not yet known.

**Tell your doctor or  
pharmacist if you notice any  
side effects from your  
medicine which are not  
mentioned here.**

---

### ***After using Medicianz MUPIROCIN Ointment***

---

#### ***Storage***

**You should keep Medicianz  
MUPIROCIN Ointment in  
the original pack below  
25°C. After first opening of  
the tube, the contents can be  
used for up to 10 days.**

**Do not freeze the ointment.**

**Do not store Medicianz  
MUPIROCIN Ointment or**

**any other medicine in the  
bathroom or near a sink.  
Do not leave it on a window  
sill or in the car.**

Heat and dampness can  
destroy some medicines.

**Keep it where children  
cannot reach it.**

A locked cupboard at least  
one-and-a-half metres above  
the ground is a good place to  
store medicines.

**Return any unused or  
expired medicine to your  
pharmacist.**

---

### ***Product description***

---

#### ***What it looks like:***

Medicianz MUPIROCIN  
Ointment is a white ointment  
supplied in 15g tubes.

#### ***Ingredients***

Medicianz MUPIROCIN  
Ointment contains the active  
ingredient Mupirocin.

Medicianz MUPIROCIN  
Ointment also contains  
polyethylene glycol (Macrogol  
400 and Macrogol 3350).

---

### ***Sponsor***

---

#### ***Marketed and distributed by:***

Medsurge Healthcare Pty Ltd

Telephone: 1300 788 261

Website: [www.medsurgehc.com](http://www.medsurgehc.com)

This leaflet was prepared in  
December 2024.