

Phentermine Lupin

Phentermine hydrochloride 15 mg and 30 mg capsules

What is in this leaflet

This leaflet answers some common questions about Phentermine Lupin. It does not contain all the available information. It does not take the place of talking to your doctor or pharmacist. If you need any more information about Phentermine Lupin please ask your doctor or pharmacist.

All medicines have risks and benefits. Your doctor has weighed the risk of you taking Phentermine Lupin against the benefits it can have for you.

If you have any concerns about taking this medicine, ask your doctor or pharmacist.

Keep this leaflet with the medicine.

You may want to read it again.

What Phentermine Lupin is used for

The name of your medicine is Phentermine Lupin.

Phentermine Lupin capsules contain the active ingredient, phentermine. Phentermine is combined with a resin which slowly releases the phentermine so the effect of Phentermine Lupin lasts all day.

Phentermine is used to reduce body weight in obese or overweight patients. Your doctor will assess whether it is suitable for you. Phentermine works by directly affecting the area of the brain that controls your appetite making you feel less hungry.

Ask your doctor if you have any questions about why this

medicine has been prescribed for you.

Phentermine should be used as a part of an overall weight management plan which should include a medically controlled diet and exercise program.

Phentermine is available only with a doctor's prescription.

Your doctor may prescribe Phentermine for another purpose. If you are not sure why you are taking this medicine ask your doctor.

Before you take Phentermine Lupin

When you must not take it

Do not take Phentermine Lupin if you have an allergy to:

- any medicine containing phentermine
- any of the ingredients listed at the end of this leaflet
- any other similar medicines used to reduce weight

Some of the symptoms of an allergic reaction may include:

- shortness of breath
- wheezing or difficulty breathing
- swelling of the face, lips, tongue, or other parts of the body
- rash, itching or hives on the skin

Do not give this medicine to a child under the age of 12 years.

Safety and effectiveness in children younger than 12 years have not been established.

Do not take this medicine if you have any of the following medical conditions:

- very high blood pressure
- high blood pressure in the lung
- problems with the valves in your heart
- disease related to blood vessels in the brain
- severe heart disease including changes in the rhythm or rate of the heart beat
- narrowing of the blood vessels due to thickening or hardening of the blood vessel wall
- overactive thyroid gland
- psychiatric illness including anorexia nervosa (loss of appetite) and depression
- glaucoma (high pressure in the eye)

Do not use this medicine:

- if you are currently taking, or have taken within the last 14 days, antidepressant medicines from the monoamine oxidase inhibitor (MAOI)
- if you have previously misused drugs or alcohol

Do not take this medicine after the expiry date printed on the pack or if the packaging is torn or shows signs of tampering.

If it has expired or is damaged, return it to your pharmacist for disposal.

If you are not sure whether you should start taking this medicine, talk to your doctor.

Before you start to take it

Tell your doctor if you have allergies to any medicines, foods, preservatives or dyes.

Tell your doctor or pharmacist if you are pregnant or plan to become pregnant or are breast-feeding.

Your doctor or pharmacist can discuss with you the risks and benefits involved.

Tell your doctor if you:

- have high blood pressure
- have heart disease such as angina (chest pain)
- have diabetes
- have epilepsy

If you have not told your doctor about any of the above, tell him/ her before you start taking Phentermine Lupin

Taking other medicines

Tell your doctor or pharmacist if you are taking any other medicines, including any that you get without a prescription from your pharmacy, supermarket or health food shop.

Some medicines and Phentermine may interfere with each other. These include:

- any medicines for weight reduction
- any medicines for depression or obsessive compulsive disorder, or other mental illness
- ergot-like medicines for migraine
- some cough and cold remedies
- thyroid hormones
- medicines for lowering high blood pressure .

These medicines may be affected by phentermine or may affect how well it works. You may need different amounts of your medicines, or you may need to take different medicines.

Your doctor and pharmacist have more information on medicines to be careful with or avoid while taking this medicine.

How to take Phentermine Lupin

Carefully follow all directions given to you by your doctor or pharmacist and never change the dose yourself.

They may differ from the information contained in this leaflet.

If you do not understand the instructions on the box, ask your doctor or pharmacist for help.

How much to take

The usual dose is one capsule every day. However, your doctor will prescribe the right dose for you.

Increasing the dose may not necessarily make you lose more weight or make you lose weight faster. However you will most likely experience more side effects. Your doctor will tell you how long you are to take Phentermine Lupin for.

How to take it

Swallow the capsules whole with plenty of water. Do not chew or open the capsules.

When to take it

Take Phentermine Lupin first thing in the morning, at breakfast time so that it does not keep you awake at night.

Taking it at the same time each day will have the best effect. It will also help you remember when to take it.

It does not matter if you take this medicine before or after food.

Phentermine Lupin is not recommended for elderly patients or for children younger than twelve years.

How long to take it

Take this medicine as long as your doctor tells you to.

If you forget to take it

If you forget to take your Phentermine Lupin dose at breakfast take it no later than lunchtime. Alternatively, skip the dose you missed and take your next dose at the normal time on the next day. If you take it any later than lunch-time you may have trouble sleeping at night.

Never take a double dose to make up for a dose you have missed.

If you have trouble remembering when to take your medicine, ask your pharmacist for some hints.

If you take too much (overdose)

Immediately telephone your doctor or the National Poisons Centre (telephone 0800 POISON or 0800 764 766) for advice, or go to Accident and Emergency at the nearest hospital, if you think that you or anyone else may have taken too much Phentermine Lupin. Do this even if there are not signs of discomfort or poisoning.

You may need urgent attention.

While you are taking it

Things you must do

If you are about to be started on any new medicine remind your doctor and pharmacist that you are taking Phentermine Lupin.

Tell any other doctors, dentists and pharmacists who treat you that you are taking this medicine.

If you are going to have surgery, tell the surgeon or

anaesthetist that you are taking this medicine.

If you become pregnant while taking this medicine, tell your doctor immediately.

If you are about to have any blood tests tell your doctor that you are taking this medicine.

It may interfere with the results of some tests.

Keep all of your doctor's appointments so that your progress can be checked.

Tell your doctor immediately if you experience sudden or rapid weight loss, difficulty breathing, chest pain, fainting, and swelling of the lower limbs and if you can not exercise as much as you usually can.

Things you must not do

**Do not take Phentermine Lupin to treat any other complaints unless your doctor tells you to.
Do not give your medicine to anyone else, even if they have the same condition as you.**

Things to be careful of

Be careful driving or operating machinery until you know how Phentermine Lupin affects you.

DO NOT drink alcohol whilst taking Phentermine Lupin.

Check with your doctor or pharmacist before you start to take any cough or cold medication.

Side effects

Tell your doctor or pharmacist as soon as possible if you do not feel well while taking Phentermine Lupin.

This medicine helps most people with a weight problem but it may have unwanted side effects in a few people. All medicines can have side effects. Sometimes they are serious, most of the time they are not. You may need medical attention if you get some of the side effects.

Do not be alarmed by the following lists of side effects. You may not experience any of them.

Ask your doctor or pharmacist to answer any questions you may have.

Tell your doctor immediately or go to Accident and Emergency at your nearest hospital if you experience:

- heart attack
- stroke
- chest pain
- myocardial infarction
- difficulty breathing

The above list includes very serious side effects. You may need urgent medical attention or hospitalisation. These side effects are rare.

Tell your doctor or pharmacist if you notice any of the following and they worry you:

- feeling more awake than usual and trouble sleeping
- your heart seems to beat faster and harder
- your blood pressure may be raised slightly
- irregular heart beats
- feeling restless
- feelings of extreme happiness followed by depression and tiredness
- nervousness
- tremor (shaking)
- headache
- dizziness
- nausea
- vomiting
- diarrhoea or constipation
- stomach cramps
- sudden or rapid weight loss
- fainting
- swelling of the lower limbs

- you cannot exercise as much as you usually can
- dry mouth or an unpleasant taste in your mouth
- difficulty passing urine
- skin rashes
- impotence
- changes in sexual drive
- swelling of the face

Tell your doctor or pharmacist if you notice anything that is making you feel unwell.

Other side effects not listed above may also occur in some people.

After taking Phentermine Lupin

Storage

Keep your capsules in the blister- foil in the carton until it is time to take them.

If you take the capsules out of the pack they may not keep well.

Keep Phentermine Lupin in a cool dry place where the temperature stays below 25°C.

Do not store Phentermine Lupin or any other medicine in the bathroom or near a sink. Do not leave them on a windowsill or in the car.

Heat and dampness can destroy some medicines.

Keep them where children cannot reach them.

A locked cupboard at least one-and-a-half metres above the ground is a good place to store medicines.

Disposal

If your doctor tells you to stop taking this medicine or the expiry date has passed, ask your pharmacist what to do with any medicine that is left over.

Product description

Date of Preparation

What it looks like

This leaflet was updated
12/12/2024

Phentermine Lupin capsules are available in two strengths in packs of 30 capsules:

15 mg - Size '3' hard gelatin capsule with white opaque cap and grey opaque body, imprinted with "PT" on cap and "15" on body in black ink and containing brown to light brown coloured spherical beads

30 mg - Size '3' hard gelatin capsule with purple cap and grey opaque body, imprinted with "PT" on cap and "30" on body in white ink and containing brown to light brown coloured spherical beads

* Not all strengths may be available.

Ingredients

Phentermine Lupin capsules contain:

- phentermine hydrochloride
- sodium polystyrene sulfonate
- magnesium stearate
- gelatin
- titanium dioxide
- iron oxide black

In addition:

Phentermine Lupin 30 mg contains:

- erythrosine
- brilliant Blue FCF

Phentermine Lupin capsules are gluten-free.

Sponsor Details

Lupin NZ Limited
c/- BDO, Level 4, Building A,
BDO Centre
4 Graham Street,
Auckland, 1010
New Zealand

Phone: +64 9 8896972