

Consumer Medicine Information (CMI) summary

The [full CMI](#) on the next page has more details. If you are worried about using this medicine, speak to your doctor or pharmacist.

1. Why am I using Yasmin?

Yasmin contains the active ingredients drospirenone and ethinylestradiol. Yasmin is a combined oral contraceptive used to prevent pregnancy. For more information, see Section [1. Why am I using Yasmin?](#) in the full CMI.

2. What should I know before I use Yasmin?

Do not use if you have ever had an allergic reaction to Yasmin or any of the ingredients listed at the end of the CMI.

Talk to your doctor if you have any other medical conditions, especially if they increase your risk of blood clots, if you have or have had blood clots, certain cancers or migraines. Tell your doctor if you take any other medicines, are pregnant or plan to become pregnant or are breastfeeding.

For more information, see Section [2. What should I know before I use Yasmin?](#) in the full CMI.

3. What if I am taking other medicines?

Some medicines may interfere with Yasmin and affect how it works. The main types of medicines that affect how well Yasmin works include certain antibiotics, antifungal medicines, HIV treatments, hepatitis C virus treatments, epilepsy medicines, medicines for high blood pressure and the heart.

A list of these medicines is in Section [3. What if I am taking other medicines?](#) in the full CMI.

4. How do I use Yasmin?

- **Swallow whole one tablet daily at around the same time each day.**
- If you vomit within 3-4 hours or have severe diarrhoea after taking a light yellow active tablet, the active ingredients may not have been completely absorbed. This is like missing a tablet. Follow the advice for missed tablets at the end of this leaflet.

More instructions can be found in Section [4. How do I use Yasmin?](#) in the full CMI.

5. What should I know while using Yasmin?

Things you should do	<ul style="list-style-type: none">• Remind any doctor, dentist or pharmacist you visit that you are using Yasmin.• Use additional barrier contraceptives (e.g. condoms) to protect yourself from STIs.• Tell your doctor if you are going to have surgery or if you plan to travel by air for more than 4 hours.• Stop taking Yasmin and call your doctor immediately if you notice signs of a possible blood clot e.g. chest pains, breathlessness or difficulty breathing, swelling, pain or tenderness in one leg, sudden dizziness, loss of balance, trouble speaking or seeing.
Things you should not do	<ul style="list-style-type: none">• Do not stop taking your medicine or change the dosage without checking with your doctor. You may become pregnant if you are not using any other contraceptive and you stop taking Yasmin, or do not take a tablet every day.
Looking after your medicine	Store it in a cool dry place away from moisture, heat or sunlight where the temperature stays below 25°C.

For more information, see Section [5. What should I know while using Yasmin?](#) in the full CMI.

6. Are there any side effects?

Less serious side effects include nausea, diarrhoea, vomiting, gut inflammation, sore throat, headache (including migraines), dizziness, mood changes, nervousness, reproductive system disorders (e.g. breast pain, unscheduled vaginal bleeding), acne, infections. Serious side effects include jaundice (yellowing skin or eyes); signs of a blood clot (e.g. breathlessness, sudden weakness/pins and needles, sudden stomach pains, fainting, unusual headaches or migraines that are worse than usual, sudden problems speaking, sudden trouble walking, pain in chest, arms or below breast bone); coughing up blood; unexplained vaginal bleeding; breast lumps. For more information, including what to do if you have any side effects, see Section [6. Are there any side effects?](#) in the full CMI.

Active ingredients: *drospirenone and ethinylestradiol*

Consumer Medicine Information (CMI)

This leaflet provides important information about using Yasmin. **You should also speak to your doctor or pharmacist if you would like further information or if you have any concerns or questions about using Yasmin.**

Where to find information in this leaflet:

- [1. Why am I using YASMIN?](#)
- [2. What should I know before I use YASMIN?](#)
- [3. What if I am taking other medicines?](#)
- [4. How do I use YASMIN?](#)
- [5. What should I know while using YASMIN?](#)
- [6. Are there any side effects?](#)
- [7. Product details](#)

1. Why am I using YASMIN?

Yasmin contains the active ingredients drospirenone and ethinylestradiol. Yasmin is a combined oral contraceptive, commonly known as a 'birth control pill' or 'the Pill'.

Yasmin is used to prevent pregnancy.

You may also experience the following benefits:

- improvement in symptoms like bloating, swelling or weight gain related to fluid retention
- more regular and lighter periods – potentially resulting in a decrease in anaemia (iron deficiency)
- a decrease in period pain.

Some conditions such as pelvic inflammatory disease, ovarian cysts, ectopic pregnancy (where the foetus is carried outside of your womb), lumpy breasts and cancer of the uterus (womb) and ovaries may be less common in women taking the Pill.

When taken correctly, it prevents you from becoming pregnant in several ways including:

- inhibiting the egg release by stopping it maturing
- changing the cervical mucus consistency, making it more difficult for the sperm to reach the egg.

When the Pill is taken by women under close observation in clinical trials, it is more than 99% effective in preventing pregnancy. However, in real life the Pill is around 92% effective. This is because pills might be missed, or taken with medicines that may interfere with their effectiveness, or may not be absorbed due to vomiting and diarrhoea.

2. What should I know before I use YASMIN?

Like all oral contraceptives, Yasmin is intended to prevent pregnancy. It does not protect against HIV infection (AIDS) and other sexually transmitted infections.

Warnings

Do not use Yasmin if:

- you are allergic to drospirenone and ethinylestradiol, or any of the ingredients listed at the end of this leaflet.
Always check the ingredients to make sure you can use this medicine.
- you are taking antiviral medicines which contain glecaprevir, pibrentasvir, sofosbuvir, velpatasvir, voxilaprevir, ombitasvir, paritaprevir, or dasabuvir, and combinations of these. These antiviral medicines are used to treat long term hepatitis C (an infectious disease that affects the liver, caused by the hepatitis C virus).
- you have or have had a blood clot in:
 - the blood vessels of the legs (deep vein thrombosis - DVT)
 - the lungs (pulmonary embolism - PE)
 - the heart (heart attack)
 - the brain (stroke)
 - other parts of the body.
- you have or are concerned about an increased risk of blood clots. You are more at risk of having a blood clot when you take the Pill. But the risk of having a blood clot when taking the Pill is less than the risk during pregnancy.
- you are concerned about an increased risk of blood clots because of age or smoking.
The risk of having a heart attack or stroke increases as you get older. It also increases if you smoke. You should stop smoking when taking the Pill, especially if you are older than 35 years of age.
- you have, or have had any blood clotting disorders such as Protein C deficiency, Protein S deficiency, Leiden Factor V mutation, Antithrombin III deficiency or other inherited blood clotting conditions
- you have, or have had:
 - a confirmed blood test showing:
 - increased levels of homocysteine
 - antiphospholipid antibodies (APLAs) e.g. anticardiolipin-antibodies and lupus anticoagulant. These may increase your risk for blood clots or miscarriages
 - major surgery after which you have not been able to move around for a period of time
 - angina (chest pain)

- mini-stroke (also known as TIA or transient ischaemic attack)
- severe kidney insufficiency or an acute failure of your kidney
- migraine, where you have also had problems with seeing, speaking or had weakness or numbness in any part of your body
- high risk of blood clots due to conditions such as diabetes with blood vessel damage, severe high blood pressure or severe high or low level of fats in your blood
- pancreatitis (an inflammation of the pancreas) associated with high levels of fatty substances in your blood
- severe liver disease and your liver function has not returned to normal
- cancer that may grow under the influence of sex hormones (e.g. of the breast or the genital organs)
- a benign or malignant liver tumour
- unexplained vaginal bleeding.

If any of these conditions appear for the first time while using the Pill, stop taking it at once and tell your doctor. In the meantime use non-hormonal (barrier) methods of contraception (such as condoms or a diaphragm).

Do not give this medicine to a child.

Check with your doctor if you:

- smoke
- you or anyone in your immediate family has had blood clots in the legs (DVT) or lungs (PE), a heart attack, a stroke, breast cancer or high cholesterol
- are overweight
- have, or have had any of the following medical conditions:
 - diabetes
 - high blood pressure
 - heart valve disorders or certain heart rhythm disorders
 - migraine
 - an increased potassium blood level (e.g. due to problems with your kidney/s) and also use diuretics or other drugs that may increase the potassium in your blood
 - cancer
 - hyperhomocysteinaemia, a condition characterised by high levels of the amino acid homocysteine in the blood
 - have any hereditary or acquired conditions that may make it more likely for you to get blood clots
 - have high cholesterol or triglycerides
 - have liver disease
 - have kidney disease
 - have high potassium in your blood
 - have jaundice (yellowing of the skin) and/or pruritus (itching of the skin) related to cholestasis (condition in which the flow of bile from the liver stops or slows)
 - have gall bladder disease

- have Crohn's disease or ulcerative colitis (chronic inflammatory bowel disease)
- have systemic lupus erythematosus (SLE – a disease affecting the skin all over the body)
- have haemolytic uraemic syndrome (HUS – a disorder of blood coagulation causing failure of the kidneys)
- have sickle cell disease
- have a condition that occurred for the first time, or worsened during pregnancy or previous use of sex hormones (e.g. hearing loss, a metabolic disease called porphyria, a skin disease called herpes gestationis, a neurological disease called Sydenham's chorea)
- have chloasma (yellowish-brown pigmentation patches on the skin, particularly of the face) – if so, avoid exposure to the sun or ultraviolet radiation
- have hereditary angio-oedema – you should see your doctor immediately if you experience symptoms of angio-oedema, such as swollen face, tongue and/or pharynx and/or difficulty swallowing, or hives together with difficulty in breathing.

If any of the above conditions appear for the first time, recur or worsen while taking Yasmin, you should tell your doctor.

During treatment, you may be at risk of developing certain side effects. It is important you understand these risks and how to monitor for them. See additional information under Section [6. Are there any side effects?](#)

Pregnancy and breastfeeding

- Check with your doctor if you are pregnant or intend to become pregnant. **Do not take this medicine if you are pregnant or think you might be pregnant.**
- **Talk to your doctor if you are breastfeeding or intend to breastfeed.** Yasmin is generally not recommended if you are breastfeeding.

3. What if I am taking other medicines?

Tell your doctor or pharmacist if you are taking any other medicines, including any medicines, vitamins or supplements that you buy without a prescription from your pharmacy, supermarket or health food shop.

Some medicines and Yasmin may interfere with each other. These include:

Medicine class	Examples
Antibiotics	rifampicin, rifabutin, macrolide antibiotics (e.g. clarithromycin, erythromycin)
Antifungal medicines	griseofulvin, ketoconazole
Medicines used to treat HIV	ritonavir or nevirapine

Medicine class	Examples
Medicines used to treat Hepatitis C Virus (HCV)	boceprevir, telaprevir, glecaprevir, pibrentasvir, ombitasvir, paritaprevir, or dasabuvir
Medicines used to treat epilepsy	phenytoin, primidone, barbiturates (e.g. phenobarbitone), carbamazepine, oxcarbazepine, topiramate, felbamate, lamotrigine
Immunosuppressants	cyclosporin
Medicines used to treat high blood pressure, chest pain or irregular heartbeats	diltiazem, verapamil, ACE inhibitors (e.g. lisinopril, perindopril), angiotensin II receptor antagonists (e.g. candesartan, irbesartan), diuretics (e.g. hydrochlorothiazide, furosemide), aldosterone antagonists (e.g. spironolactone, eplerenone)
Anti-inflammatory medicines	Etoricoxib, indomethacin
Other medicines	Tizanidine Melatonin Midazolam Theophylline St John's Wort Grapefruit juice

These medicines may be affected by Yasmin or may affect how well it works. They can:

- have an influence on the blood levels of Yasmin
- make it less effective in preventing pregnancy
- cause unexpected bleeding.

Your doctor may need to alter the dose of these medicines, or prescribe a different medicine.

You may need to use additional barrier methods of contraception (such as condoms or a diaphragm) while you are taking any of these medicines and for some time after stopping them. Your doctor will be able to tell you how long you will need to use additional contraceptive methods.

Check with your doctor or pharmacist if you are not sure about what medicines, vitamins or supplements you are taking and if these affect Yasmin.

4. How do I use YASMIN?

How much to take

- Take one tablet daily at about the same time every day. You must take Yasmin every day regardless of how often you have sex. This will also help you remember when to take it
- **Swallow the tablet whole with a full glass of water.** It does not matter if you take it before or after food.
- If you vomit within 3-4 hours or have severe diarrhoea after taking a light yellow active tablet, the active ingredients may not have been completely absorbed.

This is like missing a tablet. Follow the advice for missed tablets.

How to take Yasmin

- Each blister pack is marked with the day of the week.
- Take your first light yellow (active) tablet from the green area on the blister pack corresponding to the day of the week.
- Follow the direction of the arrows on the blister pack until all the tablets have been taken.
- A period should begin 2-3 days after starting to take the white inactive tablets and may not have finished before the next pack is started.
- Always start a new blister pack on the same day of the week as your previous pack.

Taking Yasmin for the first time

- If you are starting Yasmin after a natural cycle, and you have not used a hormonal contraceptive in the past month, start on the first day of your period, i.e. on the first day of your menstrual bleeding.
- You may also start on days 2-5 of your period, but in that case make sure you also use additional barrier contraceptive precautions (e.g. condom) for the first 7 days of tablet-taking.
- Your doctor will advise you when to start if you:
 - are taking Yasmin after having a baby
 - have had a miscarriage or an abortion.

Switching from another contraceptive

Type of contraceptive	How do I change over to Yasmin?	Will I need additional contraception?
Combined oral contraceptive	Start taking Yasmin on the day after taking the last active tablet in your previous pill pack. Bleeding may not occur until the end of the first pack of Yasmin. You can also switch to Yasmin after taking one or more inactive tablets in your previous pack. But no later than the day after taking the last inactive tablet	No
Vaginal ring	Start Yasmin on the day of removal of the vaginal ring or at the latest when the next application would have been due	No
Progesterone only pill (minipill)	Stop taking the minipill on any day and start taking	You must also use additional barrier

	Yasmin at the same time the day after you took your last minipill	contraceptive precautions (e.g. condoms or a diaphragm) for the first 7 days of tablet-taking when having intercourse
Progestogen-only injection, implant or intrauterine system (IUS)	Start taking Yasmin when your next injection is due, or on the day that your implant or IUS is removed	

Stopping Yasmin

You can stop taking Yasmin at any time. If you are considering becoming pregnant, it is recommended that you begin taking a vitamin supplement containing folic acid. It is best that you start taking folic acid tablets before you stop taking Yasmin and not stop until your doctor advises this. Ask your doctor or pharmacist about suitable supplements. It is both safe and recommended that you take folic acid during pregnancy.

If you forget to use Yasmin

For Yasmin to be most effective, light yellow active tablets need to be taken uninterrupted for 7 days.

If you miss a tablet and take the missed tablet within 12 hours of missing it, you should still be protected against pregnancy. If you are more than 12 hours late follow these detailed instructions:

- **If you have been taking the light yellow active tablets for 7 uninterrupted days and miss a light yellow active tablet**, take the missed tablet as soon as you remember, then go back to taking your medicine as you would normally, even if this means taking two tablets in one day. You will not need to use additional barrier contraceptive precautions.
 - The chance of pregnancy after missing a light yellow active tablet depends on when you missed the tablet. **There is a higher risk of becoming pregnant if you miss a tablet at the beginning or end of a pack.**
 - If after taking your missed tablet you have less than 7 days of light yellow active tablets left in a row, you should finish the active tablets in your pack but skip the white inactive tablets and start a new pack. This is the best way to maintain contraceptive protection. However, you may not have a period until the end of the light yellow active tablets of the second pack. You may have spotting or breakthrough bleeding on tablet-taking days.
- **If you have been taking the light yellow active tablets for less than 7 days and miss a light yellow active tablet**, take the missed tablet as soon as you remember, then go back to taking your medicine as you would normally, even if this means taking two tablets in one day.
 - **You must also use additional barrier contraceptive precautions (e.g. condoms or a diaphragm) for the next 7 days.**

- If you have had sexual intercourse during that time, there is a possibility of pregnancy and you may need emergency contraception.
- **If you forget to take more than one light yellow active tablet**, seek advice from your doctor or pharmacist about what to do.
 - If you have had sexual intercourse in the week before missing your tablets, there is a possibility of becoming pregnant.
- **If you miss a white inactive tablet, you do not need to take them later because they do not contain any active ingredients.** However, it is important that you discard the missed white tablet(s) to make sure that the number of days between taking active tablets is not increased as this would increase the risk of pregnancy. Continue with the next tablet at the usual time.

Please see the [diagram](#) at the end of this leaflet for "Summary of advice if you missed a light yellow active tablet more than 12 hours ago".

If you use too much Yasmin

If you think that you have used too much Yasmin, you may need urgent medical attention.

You should immediately:

- phone the Poisons Information Centre
(by calling **13 11 26** in Australia or **0800 764 766** in New Zealand), or
- contact your doctor, or
- go to the Emergency Department at your nearest hospital.

You should do this even if there are no signs of discomfort or poisoning.

If you take several light yellow active tablets at once, you may feel sick or vomit or may bleed from the vagina. Even girls who have not yet started to menstruate but have accidentally taken this medicine may experience such bleeding.

5. What should I know while using YASMIN?

Things you should do

- Tell any doctors, dentists and pharmacists who treat you that you are taking this medicine.
- Yasmin will not protect you from HIV-AIDS or any other Sexually Transmitted Infections (STIs), such as chlamydia, genital herpes, genital warts, gonorrhoea, hepatitis B, human papilloma virus and syphilis.
To protect yourself from STIs, you will need to use additional barrier contraceptives (e.g. condoms).
- If you are about to have any blood tests, tell your doctor that you are taking this medicine. It may interfere with the results of some tests.
- **Have regular check-ups with your doctor.** When you are taking the Pill, your doctor will tell you to return for regular check-ups, including getting a Cervical Screening Test. Your doctor will advise how often you

need a Cervical Screening Test. A Cervical Screening Test can detect abnormal cells lining the cervix. Sometimes abnormal cells can progress to cancer.

- If you are about to start on any new medicine, remind your doctor and pharmacist that you are taking Yasmin
- **If you are going to have surgery, tell the surgeon or anaesthetist beforehand that you are taking this medicine.** The risk of having blood clots is temporarily increased as a result of major surgery, any surgery to the legs or pelvis, neurosurgery or major trauma. In women who take Yasmin, the risk may be higher.
- In women at risk of prolonged immobilisation (including major surgery, any surgery to the legs or pelvis, neurosurgery, or major trauma), your doctor may tell you to stop taking (in the case of elective surgery at least four weeks in advance) and not resume until two weeks after complete remobilisation. Another method of contraception should be used to avoid unintentional pregnancy. Your doctor may prescribe other treatment (e.g. treatment for blood clots) if Yasmin has not been discontinued in advance.
- **Consult your doctor if you plan to air travel for greater than 4 hours.** Other risk factors for blood clotting include temporary immobilisation including air travel of greater than 4 hours, particularly in women with other risk factors.
- **Consult your doctor if you develop high blood pressure while taking Yasmin** – you may be told to stop taking it.
- If you become pregnant while taking this medicine, tell your doctor immediately.

Call your doctor straight away if you notice:

- one-sided swelling of the leg and/or foot or along a vein in the leg
- pain or tenderness in the leg which may be felt only when standing or walking
- increased warmth in the affected leg; red or discoloured skin on the leg
- sudden onset of unexplained shortness of breath or rapid breathing
- sudden coughing or coughing up of blood
- sharp chest pain or sudden severe pain in the chest which may increase with deep breathing
- severe light headedness or dizziness
- rapid or irregular heartbeat
- sudden pain, swelling and slight blue discoloration of an extremity
- sudden numbness or weakness of the face, arm or leg, especially on one side of the body
- sudden trouble walking, dizziness, loss of balance or coordination
- sudden confusion, slurred speech or aphasia; sudden partial or complete loss of vision, double vision, painless blurring of vision which can progress to loss of vision
- sudden, severe or prolonged headache with no known cause
- loss of consciousness or fainting with or without seizure

- pain, discomfort, pressure, heaviness, sensation of squeezing or fullness in the chest arm, or below the breastbone
- discomfort radiating to the back, jaw, throat, arm, stomach
- feeling of being full, having indigestion or choking
- sweating, nausea, vomiting
- extreme weakness and anxiety.

Things you should not do

- Do not take Yasmin to treat any other conditions, unless your doctor tells you to.
- Do not give your medicine to anyone else.
- **Do not stop taking your medicine or change the dosage without checking with your doctor.** You may become pregnant if you are not using any other contraceptive and you stop taking Yasmin, or do not take a tablet every day.

Bleeding and Missed Periods

- **If you have unexpected bleeding and it continues, becomes heavy, or occurs again, tell your doctor.**
- When taking these tablets for the first few months, you can have irregular vaginal bleeding (spotting or breakthrough bleeding) between your periods. You may need to use sanitary products, but continue to take your tablets as normal. Irregular vaginal bleeding usually stops once your body has adjusted to the Pill, usually after about 3 months.
- If you have missed a period, but you have taken all your tablets, it is very unlikely that you are pregnant, as long as:
 - you have taken the light yellow active tablets at the right time
 - you have not been taking medicine(s) that may interfere with Yasmin
 - you have not vomited or had severe diarrhoea during this cycle.
- If this is so, continue to take Yasmin as usual. If you have any concerns consult your doctor or pharmacist.
- **If you miss your period twice in a row, you may be pregnant even if you have taken the Pill correctly. Stop taking Yasmin and seek advice from your doctor. You must use a non-hormonal method of contraception (such as condoms or a diaphragm) until your doctor rules out pregnancy.**

Driving or using machines

Be careful before you drive or use any machines or tools until you know how Yasmin affects you.

Looking after your medicine

Follow the instructions in the carton on how to take care of your medicine properly.

Store it in a cool dry place away from moisture, heat or sunlight where the temperature stays below 25°C.

Do NOT store it in the bathroom or near a sink, or in the car or on window sills.

Keep it where young children cannot reach it.

Getting rid of any unwanted medicine

If you no longer need to use this medicine or it is out of date, take it to any pharmacy for safe disposal.

Do not use this medicine after the expiry date.

6. Are there any side effects?

All medicines can have side effects. If you do experience any side effects, most of them are minor and temporary. However, some side effects may need medical attention.

See the information below and, if you need to, ask your doctor or pharmacist if you have any further questions about side effects.

Less serious side effects

Less serious side effects	What to do
Gastrointestinal system-related <ul style="list-style-type: none">nauseadiarrhoeavomitinginflammation of the stomach and intestinessore throat and discomfort when swallowing Nervous system-related <ul style="list-style-type: none">headache, including migrainesdizziness Mood-related <ul style="list-style-type: none">mood changes, including depressionnervousness Reproductive system-related <ul style="list-style-type: none">menstrual disordersbreast painabnormal vaginal discharge Skin-related <ul style="list-style-type: none">acne Infections <ul style="list-style-type: none">inflammation of the bladder due to urinary tract infectionvaginal yeast infectionsinus infection	Speak to your doctor if you have any of these less serious side effects and they worry you.

Serious side effects

Serious side effects	What to do
Blood clot-related: <ul style="list-style-type: none">pain in the chest, arm or below the breastbonepain or discomfort that goes to your back	Call your doctor straight away, or go straight to the Emergency

<ul style="list-style-type: none">breathlessness and/or difficulty breathingswelling, pain or tenderness of one legsudden weakness, numbness or bad 'pins and needles' of the face, arm or leg, especially on one side of the bodysudden trouble walking, dizziness, loss of balance or coordinationsevere, sudden stomach painsa fainting attack, or you collapseunusual headaches or migraines that are worse than usualsudden problems with your speech, understanding or eyesight Liver-related <ul style="list-style-type: none">jaundice (yellowing skin or yellowing eyes) Bleeding-related <ul style="list-style-type: none">coughing up blood Reproductive system-related <ul style="list-style-type: none">breast lumpsunexplained vaginal bleeding	Department at your nearest hospital if you notice any of these serious side effects.
--	---

Tell your doctor or pharmacist if you notice anything else that may be making you feel unwell.

Other side effects not listed here may occur in some people.

Blood clots and the Pill

Blood clots are rare. Very occasionally blood clots may cause serious permanent disability and may even be fatal.

Blood clots can form, travel and block different blood vessels in the body. They can occur in the deep veins of the legs (causing deep vein thrombosis), the clot can travel and cause a blockage in the lungs (called pulmonary embolism). Blood clots can occur in the blood vessels of the heart (causing a heart attack) or the brain (causing a stroke).

All combined oral contraceptive pills, including Yasmin will increase the risk of having a blood clot. However, blood clots can form whether or not you are taking the pill, including in pregnancy. The risk of having a blood clot when taking Yasmin is less than the risk of having a blood clot during pregnancy.

The risk of a blood clot is highest during the first year of taking the Pill for the first time, or when re-starting after having a break from the Pill for 4 weeks or more.

Stop taking Yasmin and call your doctor straight away, or go straight to the Emergency Department at your nearest hospital if you notice signs of a possible blood clot.

To prevent pregnancy, you must also use additional barrier contraceptive precautions (e.g. condoms or a diaphragm).

Cancer and the Pill

- Breast cancer has been diagnosed slightly more often in women who take the Pill than in women of the same age who do not take the Pill.
- This slight increase in the numbers of breast cancer diagnoses gradually disappears during the course of the 10 years after women stop taking the Pill.
- It is not known whether the difference is caused by the Pill. It may be that these women were examined more often, so that the breast cancer was noticed earlier.
- **It is important that you check your breasts regularly and contact your doctor if you feel any lumps.**
- In rare cases benign liver tumours and, even more rarely, malignant liver tumours have been reported in users of the Pill. These tumours may lead to internal bleeding.
- **Contact your doctor immediately if you have severe pain in your abdomen.**
- Cervical cancer has been reported to occur more often in women who have been taking the Pill for a long time. This finding may not be caused by the Pill, but may be related to sexual behaviour and other factors.

Reporting side effects

After you have received medical advice for any side effects you experience, you can report side effects to the Therapeutic Goods Administration online at www.tga.gov.au/reporting-problems or in New Zealand at <https://pophealth.my.site.com/carmreportnz/s/>. By reporting side effects, you can help provide more information on the safety of this medicine.

Always make sure you speak to your doctor or pharmacist before you decide to stop taking any of your medicines.

7. Product details

This medicine is only available with a doctor's prescription.

What Yasmin contains

Active ingredients (main ingredients in the light yellow tablets)	<ul style="list-style-type: none">• 3 mg drospirenone• 30 microgram ethinylestradiol
--	---

Other ingredients (inactive ingredients)	Light yellow tablet <ul style="list-style-type: none">• lactose monohydrate• maize starch• pregelatinised maize starch• povidone• magnesium stearate• hypromellose• macrogol 6000• purified talc• titanium dioxide• iron oxide yellow White tablet <ul style="list-style-type: none">• lactose monohydrate• microcrystalline cellulose• magnesium stearate• hypromellose• purified talc• titanium dioxide
Potential allergens	lactose monohydrate

Do not take this medicine if you are allergic to any of these ingredients.

What Yasmin looks like

Yasmin active tablets are light yellow round tablets with 'DO' marked on one side in a regular hexagon.

Yasmin inactive tablets are round white tablets marked on one side with the letters 'DP' in a regular hexagon.

Yasmin comes in a box containing either 1 or 3 blister packs. Each blister pack contains 21 light yellow active tablets and 7 white inactive tablets. (Aust R 226238)

Who distributes Yasmin

Bayer Australia Ltd
ABN 22 000 138 714
875 Pacific Highway
Pymble, NSW 2073

Bayer New Zealand
Hillcrest North Shore
Auckland 0627
New Zealand

See TGA website (www.ebs.tga.gov.au) for latest Australian Consumer Medicine Information.

See MEDSAFE website (www.medsafe.govt.nz) for latest New Zealand Consumer Medicine Information.

® Registered Trademark of the Bayer Group, Germany

© Bayer Australia Ltd
All rights reserved.

This leaflet was prepared in August 2024.

Summary of advice if you missed a light yellow active tablet more than 12 hours ago

Before missing your tablet, did you take light yellow active tablets for the previous 7 days?	No	Did you have sex in the 7 days before missing the tablet?	→	<p>No</p> <p>Take the tablet missed AND use extra barrier precaution for 7 days. If there are fewer than 7 light yellow active tablets left in the pack, finish the active tablets and go straight to the light yellow active tablets of the next pack. This means you skip the white inactive tablets</p> <p>Yes</p> <p>See your Doctor or Pharmacist for advice</p>
	Yes	Does your pack still have 7 active light yellow tablets in a row to follow?	→	<p>No</p> <p>Take the tablet missed AND complete taking the light yellow active tablets. Skip the white inactive tablets. Start your next pack with the light yellow active tablets</p> <p>Yes</p> <p>Take the tablet missed AND complete the pack as normal</p>